
Native Grass Seeds



Slender Wheatgrass

Elymus trachycaulus

- somewhat drought tolerant
- can withstand 21-63 days flooding
- widely used for revegetation



Percy Folkard 2013

Fringed Brome

Bromus ciliatus L.

- effective for erosion control
- seeds provide food for small mammals and birds




Bryan Kelly-McArthur 2021

Sloughgrass

Beckmannia

- grows in wet meadows and along pond edges
- used in erosion control and land reclamation



Ryan Batten 2014

Fowl Bluegrass

Poa palustris

- grows in wet meadows and riparian sites
- used for erosion control and revegetation




Jaime Fenneman 2006

Streambank Wheatgrass

Elymus lanceolatus subsp. riparius

drought tolerant

used for stabilization of disturbed soils




Elymus lanceolatus ssp. *lanceolatus* The Illustrated Flora of British Columbia

Northern Wheatgrass (Thickspike Wheatgrass)

Elymus lanceolatus

- food for deer in spring
- used for stabilization of disturbed soils



Loren St. John

Junegrass

Koeleria macrantha

- food for deer in spring
- used for stabilization of disturbed soils




Adolf Ceska 2008

Tufted Hairgrass

Deschampsia cespitosa (L.) P. Beauv.

- well suited to moist meadows
- used for shoreline stabilization and revegetation



Harry Hill 2017